GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

Fundamentals of Schiff-Base Ligands and Their Transition-Metal Complexes

Pushpa, Shiv Brat Singh
Research Scholar, Department of Chemistry, Surendranagar University,
Shree Pandit Nathulalji Vyas Campus, Surendranagar, Gujarat, India

Abstract

Schiff bases are an important family of compounds formed through condensation reactions between primary amines and aldehydes or ketones. Their imine nitrogen, together with additional donor atoms introduced by suitable substituents, enables coordination to transition-metal ions. Semicarbazones and thiosemicarbazones are particularly useful derivatives because they offer nitrogen-, oxygen- and/or sulfur-containing donor environments. This review introduces the fundamental chemistry relevant to the preparation and study of such ligands and their cobalt, nickel and copper complexes. It summarises general synthetic principles, coordination behaviour, complementary characterisation methods and the rationale for selected in-vitro antioxidant and antibacterial assays. Emphasis is placed on sound interpretation: spectral shifts may support coordination but do not independently establish a complete structure and preliminary assay responses should not be described as therapeutic efficacy. The discussion provides a conceptual framework for understanding thesis research involving semicarbazone and thiosemicarbazone ligands derived from aromatic and cyclic carbonyl compounds.

Keywords: Schiff base; semicarbazone; thiosemicarbazone; transition-metal complex; coordination chemistry; spectroscopy; antioxidant assay; antibacterial screening.

1. Introduction

Schiff bases, commonly described as imine compounds, are characterised by a carbon–nitrogen double bond (C=N). They are generally prepared by reacting a primary amine with an aldehyde or ketone, often under conditions that facilitate removal of water. The ease of synthesis and the ability to vary both the carbonyl precursor and the amine component make these compounds adaptable ligand platforms. Their coordination chemistry and reported applications have been widely reviewed in the chemical literature [1–4].

GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

In coordination chemistry, a ligand is an ion or molecule that donates one or more electron pairs to a central metal ion. Schiff-base ligands can bind through the azomethine nitrogen and, where present, additional donor atoms such as oxygen or sulfur. Semicarbazones and thiosemicarbazones are relevant examples in which the terminal functional groups provide further opportunities for metal binding.

2. Formation and classes of Schiff-base ligands

The usual condensation pathway involves nucleophilic addition of the amine to the carbonyl carbon, formation of a carbinolamine-type intermediate, proton transfers and elimination of water to generate the C=N linkage. Reaction conditions, solvent, pH, temperature and the structure of the reactants can influence conversion, product distribution and isolation.



Figure 1. Simplified conceptual pathway for Schiff-base formation.

2.1 Semicarbazones

Semicarbazones are obtained by condensation of semicarbazide with an aldehyde or ketone. They contain an imine-related nitrogen and a carbonyl-containing urea-like fragment. Depending on substitution and tautomeric state, the ligand may provide more than one potential donor site. The actual donor set must be determined for each compound rather than assumed from its class.

2.2 Thiosemicarbazones

Thiosemicarbazones are formed from carbonyl compounds and thiosemicarbazide. Their sulfur-containing group introduces additional coordination possibilities. Thione–thiol tautomerism and

GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

deprotonation can affect the preferred binding mode. Many reported thiosemicarbazone complexes show diverse structures and biological responses, but these properties depend on ligand substitution, metal identity, oxidation state, geometry and experimental conditions [2,3].

Ligand family	Typical distinguishing feature	Potential donor atoms*
Semicarbazone	O-containing semicarbazide-derived group	N; O and/or other sites depending on structure
Thiosemicarbazone	S-containing thiosemicarbazide-derived group	N; S and sometimes additional donor sites

***Potential donor atoms are illustrative; actual coordination must be confirmed experimentally.**

3. Coordination of Schiff-base ligands to transition metals

Transition-metal ions accept electron pairs from ligand donor atoms to form coordinate bonds. The number and arrangement of donor atoms around a metal centre define its coordination number and geometry. Common geometries include tetrahedral, square-planar and octahedral arrangements, although distortions and other geometries are also possible. The preferred structure depends on the metal ion, oxidation state, electronic configuration, ligand denticity, steric effects, counter-ions and solvent.



GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

Figure 2. Generic donor–metal coordination concept; not a proposed structure for any specific thesis compound.

3.1 Cobalt, nickel and copper

Metal ion	General coordination-chemistry considerations	Characterisation considerations
Co(II)	May show several coordination numbers and geometries; magnetic behaviour can be informative.	Magnetic susceptibility and electronic spectra can help evaluate possible environments.
Ni(II)	Can adopt square-planar, tetrahedral or octahedral environments depending on ligand field and conditions.	UV–Visible spectra and magnetic data are useful alongside other evidence.
Cu(II)	Often exhibits Jahn–Teller effects and distorted coordination environments.	Electronic spectra, EPR where available and crystallography can help resolve geometry.

These are general trends, not assignments. A particular geometry should not be declared solely from the identity of the metal or a single absorption band. Elemental analysis, magnetic measurements, mass spectrometry, EPR and single-crystal X-ray diffraction can provide additional evidence where available.

4. General approaches to synthesis

Ligand synthesis commonly uses a carbonyl compound and the appropriate nitrogen-containing reagent in a suitable solvent. Acid or base catalysis may be used depending on the substrate and desired product. Isolation can involve cooling, filtration, washing and drying. Metal-complex synthesis often involves combining a ligand solution with a metal salt solution under controlled stoichiometry, solvent and temperature. Reflux is frequently used to promote reaction, followed by isolation and washing of the solid product.

Reproducible reporting should state reagent identity and purity, molar quantities, solvent volumes, pH where relevant, temperature, reaction time, work-up procedure, isolated mass, calculated yield and storage conditions. The proposed empirical formula and metal-to-ligand ratio must be consistent with analytical evidence rather than inferred only from the reagent feed ratio.

GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

5. Characterisation: combining complementary evidence

No single analytical method usually establishes every aspect of a coordination compound. Each technique answers a different question and the strongest structural interpretation comes from mutually consistent evidence.



Figure 3. Complementary information obtained from common analytical methods.

Technique	Typical purpose	Important caution
TLC	Monitor reaction progress and compare spot patterns.	A single spot does not prove structural identity or absolute purity.
IR spectroscopy	Identify functional groups and examine changes in C=N, C=O, N–H or C=S-related regions.	Band shifts support an interpretation but assignments need full spectra and careful comparison.
NMR spectroscopy	Investigate chemical environments and ligand structure.	Paramagnetic metal complexes may produce broadened or shifted signals; applicability varies.
UV–Visible spectroscopy	Study ligand-centred and metal-related electronic transitions.	Assignments should be supported by metal configuration and other measurements.
HPLC	Separate components and assess chromatographic purity under stated conditions.	Retention time alone is not a structural identification.

GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

6. Biological screening: what the assays measure

Antioxidant-related and antibacterial assays are frequently used as preliminary screens for newly prepared compounds. Different assays probe different chemical or biological processes; their percentage values should not be directly compared as if they measured the same property.



Figure 4. Overview of distinct in-vitro screening endpoints.

6.1 DPPH radical-scavenging assay

The DPPH assay uses a stable coloured radical. A decrease in absorbance after addition of a test sample is interpreted as radical-scavenging response under the selected conditions. Results depend on solvent, reaction time, sample concentration, light exposure and calculation method. A suitable positive control and sample blanks are important.

6.2 Nitric-oxide-related assay

Common nitric oxide scavenging protocols generate nitric oxide from a donor compound and estimate nitrite formation using a colour-forming reagent system. The response is influenced by incubation time, temperature, pH and reagent concentrations. The exact protocol and calibration approach should be reported.

6.3 Hydrogen-peroxide scavenging assay

This assay estimates the reduction in hydrogen peroxide under defined conditions, often using UV absorbance. Because compounds may absorb or scatter light themselves, sample blanks and appropriate controls are required to avoid misleading calculations.

GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

6.4 Agar-well diffusion antibacterial assay

In agar-well diffusion, a test solution is placed in a well cut into inoculated agar and the zone of growth inhibition is measured after incubation. Zone size depends not only on antimicrobial effect but also on diffusion, solubility, concentration, agar composition and inoculum density. Reports should identify the organism and strain, inoculum standard, medium, well diameter, test concentration, incubation conditions, controls and measured zone diameters. MIC and MBC measurements can strengthen the evaluation.

7. Interpreting structure–property relationships

Changes in ligand donor atoms, substituents, metal identity and coordination geometry can alter charge distribution, stability, solubility and redox behaviour. These changes may influence assay responses, but a simple claim that metal coordination always improves biological activity is not justified. Reliable structure–activity discussion requires verified structures, comparable concentrations, replicated measurements and statistical analysis. Potential mechanisms should be presented as hypotheses unless directly tested.

Likewise, in-vitro antioxidant or antibacterial activity does not establish safety, selectivity, bioavailability or clinical usefulness. Further studies may include MIC/MBC testing, cytotoxicity against appropriate mammalian cell lines, mechanistic experiments, stability studies and, where justified, in-vivo evaluation.

8. Relevance to thesis-based research

The thesis framework underlying this companion article considers four ligand systems: PS and PTS derived from 3-chloro-3-phenyl-2-propenal and CS and CTS derived from cyclohex-2-enone. PS and CS are described as semicarbazone derivatives, while PTS and CTS are described as thiosemicarbazone derivatives. The ligand families are considered with Co(II), Ni(II) and Cu(II), using chloride- and nitrate-derived preparations, followed by chromatographic and spectroscopic characterisation and selected biological screening. These features provide a practical case study for the fundamental concepts reviewed above.

This article deliberately focuses on the underlying chemistry and the interpretation of methods rather than repeating the numerical results of the separate research article. Before submission, all thesis-

GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

specific ligand names, formulae, spectra, reagent quantities and protocols should be checked against original laboratory records and the final thesis.

9. Research gaps and future directions

- Confirm molecular structures and metal-to-ligand stoichiometry using multiple independent analytical methods.
- Where possible, obtain single-crystal X-ray diffraction or complementary structural evidence for definitive geometry assignments.
- Report full spectra, peak tables, elemental analysis, magnetic data and mass-spectrometric evidence.
- Standardise biological assay concentrations, controls, replicates, bacterial strain identity and statistical treatment.
- Add MIC/MBC and cytotoxicity studies before making broader biological claims.
- Explore greener synthesis approaches, solvent minimisation and reproducible yield optimisation.

10. Conclusion

Schiff bases, including semicarbazones and thiosemicarbazones, offer versatile platforms for transition-metal coordination chemistry. Their study connects organic synthesis, inorganic coordination, spectroscopy and preliminary biological evaluation. A robust research report must distinguish proposed structures from experimentally established ones, use complementary analytical evidence and interpret biological assays within their limitations. The fundamentals summarised here provide a framework for understanding and critically evaluating thesis-based studies of cobalt, nickel and copper complexes.

References

1. More, M. S., Joshi, P. G., Mishra, Y. K. and Khanna, P. K. (2019). Metal complexes driven from Schiff bases and semicarbazones for biomedical and allied applications: A review. Materials Today Chemistry, 14, 100195.
2. Chugh, V., Mohan, C., Meena, P. L., Mangla, D. and Zaidi, N. (2026). Synthesis, characterization and biological applications of semicarbazide and thiosemicarbazide derived Schiff base complexes. Discover Materials.

GYANBODH
An International Multidisciplinary Peer Reviewed Journal
ISSN:3048-9792
Volume: 3
Issue: 5
September-October: 2026

3. Marinova, P. E. and Tamahkyarova, K. D. (2025). Synthesis, investigation, biological evaluation and application of coordination compounds with Schiff base—A review. Compounds, 5(2), 14.
4. Malav, R. and Ray, S. (2025). Recent advances in the synthesis and versatile applications of transition metal complexes featuring Schiff base ligands. RSC Advances, 15, 22889–22914.
5. Pushpa and Shiv Brat Singh. (2024). Study of biological activities and characterisation of cobalt, nickel and copper complexes synthesised from Schiff base derivatives derived from 3-chloro-3-phenyl-2-propenal and cyclohex-2-enone. AYUDH: International Peer-Reviewed Refereed Journal, 6, 31–36.
6. Pushpa and Shiv Brat Singh. (2026). Synthesis, Spectral Characterization and Antioxidant–Antimicrobial Evaluation of Zinc(II) Schiff Base Complexes Derived from α,β-Unsaturated Carbonyl Compounds. Research Genius 11(21): 501-502.